



spondylocarpotarsal synostosis syndrome

Spondylocarpotarsal synostosis syndrome is a disorder that affects the development of bones throughout the body. Newborns with this disorder are of approximately normal length, but impaired growth of the trunk results in short stature over time. The bones of the spine (vertebrae) are misshapen and abnormally joined together (fused). The vertebral abnormalities may result in an abnormally curved lower back (lordosis) and a spine that curves to the side (scoliosis).

Affected individuals also have abnormalities of the wrist (carpal) and ankle (tarsal) bones and inward- and upward-turning feet (clubfeet). Characteristic facial features include a round face, a large forehead (frontal bossing), and nostrils that open to the front rather than downward (anteverted nares).

Some people with spondylocarpotarsal synostosis syndrome have an opening in the roof of the mouth (a cleft palate), hearing loss, thin tooth enamel, flat feet, or an unusually large range of joint movement (hypermobility). Individuals with this disorder can survive into adulthood. Intelligence is generally unaffected, although mild developmental delay has been reported in some affected individuals.

Frequency

Spondylocarpotarsal synostosis syndrome is a rare disorder; its prevalence is unknown. At least 25 affected individuals have been identified.

Genetic Changes

Mutations in the *FLNB* gene cause spondylocarpotarsal synostosis syndrome. The *FLNB* gene provides instructions for making a protein called filamin B. This protein helps build the network of protein filaments (cytoskeleton) that gives structure to cells and allows them to change shape and move. Filamin B attaches (binds) to another protein called actin and helps the actin to form the branching network of filaments that makes up the cytoskeleton. It also links actin to many other proteins to perform various functions within the cell, including the cell signaling that helps determine how the cytoskeleton will change as tissues grow and take shape during development.

Filamin B is especially important in the development of the skeleton before birth. It is active (expressed) in the cell membranes of cartilage-forming cells (chondrocytes). Cartilage is a tough, flexible tissue that makes up much of the skeleton during early development. Most cartilage is later converted to bone (a process called ossification), except for the cartilage that continues to cover and protect the ends of bones and is present in the nose, airways (trachea and bronchi), and external ears. Filamin

B appears to be important for normal cell growth and division (proliferation) and maturation (differentiation) of chondrocytes and for the ossification of cartilage.

FLNB gene mutations that cause spondylocarpotarsal synostosis syndrome result in the production of an abnormally short filamin B protein that is unstable and breaks down rapidly. Loss of the filamin B protein appears to result in out-of-place (ectopic) ossification, resulting in fusion of the bones in the spine, wrists, and ankles and other signs and symptoms of spondylocarpotarsal synostosis syndrome.

A few individuals who have been diagnosed with spondylocarpotarsal synostosis syndrome do not have mutations in the *FLNB* gene. In these cases, the genetic cause of the disorder is unknown.

Inheritance Pattern

Spondylocarpotarsal synostosis syndrome caused by *FLNB* gene mutations is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

In a few individuals with signs and symptoms similar to those of spondylocarpotarsal synostosis syndrome but without *FLNB* gene mutations, the condition appears to have been inherited in an autosomal dominant pattern. Autosomal dominant means one copy of the altered gene in each cell is sufficient to cause the disorder.

Other Names for This Condition

- congenital scoliosis with unilateral unsegmented bar
- congenital synspondylism
- SCT
- SCT syndrome
- spondylocarpotarsal syndrome
- vertebral fusion with carpal coalition

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Spondylocarpotarsal synostosis syndrome
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1848934/>

Other Diagnosis and Management Resources

- GeneReview: *FLNB*-Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK2534>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Health Topic: Bone Diseases
<https://medlineplus.gov/bonediseases.html>
- Health Topic: Scoliosis
<https://medlineplus.gov/scoliosis.html>

Genetic and Rare Diseases Information Center

- Spondylocarpotarsal synostosis syndrome
<https://rarediseases.info.nih.gov/diseases/4974/spondylocarpotarsal-synostosis-syndrome>

Educational Resources

- Disease InfoSearch: Spondylocarpotarsal synostosis syndrome
<http://www.diseaseinfosearch.org/Spondylocarpotarsal+synostosis+syndrome/6801>
- MalaCards: spondylocarpotarsal synostosis syndrome
http://www.malacards.org/card/spondylocarpotarsal_synostosis_syndrome
- Orphanet: Spondylocarpotarsal synostosis
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=3275

Patient Support and Advocacy Resources

- Little People of America
<http://www.lpaonline.org/>
- The MAGIC Foundation for Children's Growth
<https://www.magicfoundation.org/>

GeneReviews

- FLNB-Related Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK2534>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28spondylocarpotarsal+synostosis+syndrome%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- SPONDYLOCARPOTARSAL SYNOSTOSIS SYNDROME
<http://omim.org/entry/272460>

Sources for This Summary

- Isidor B, Cormier-Daire V, Le Merrer M, Lefrancois T, Hamel A, Le Caignec C, David A, Jacquemont S. Autosomal dominant spondylocarpotarsal synostosis syndrome: phenotypic homogeneity and genetic heterogeneity. *Am J Med Genet A*. 2008 Jun 15;146A(12):1593-7. doi: 10.1002/ajmg.a.32217.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/18470895>
- Krakow D, Robertson SP, King LM, Morgan T, Sebald ET, Bertolotto C, Wachsmann-Hogiu S, Acuna D, Shapiro SS, Takafuta T, Aftimos S, Kim CA, Firth H, Steiner CE, Cormier-Daire V, Superti-Furga A, Bonafe L, Graham JM Jr, Grix A, Bacino CA, Allanson J, Bialer MG, Lachman RS, Rimoin DL, Cohn DH. Mutations in the gene encoding filamin B disrupt vertebral segmentation, joint formation and skeletogenesis. *Nat Genet*. 2004 Apr;36(4):405-10. Epub 2004 Feb 29.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14991055>
- Sawyer GM, Clark AR, Robertson SP, Sutherland-Smith AJ. Disease-associated substitutions in the filamin B actin binding domain confer enhanced actin binding affinity in the absence of major structural disturbance: Insights from the crystal structures of filamin B actin binding domains. *J Mol Biol*. 2009 Jul 31;390(5):1030-47. doi: 10.1016/j.jmb.2009.06.009. Epub 2009 Jun 6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19505475>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/spondylocarpotarsal-synostosis-syndrome>

Reviewed: September 2011

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services